

# Production protocol

## Omeprazole suspension

(2-5 mg/mL) with Liquifine® Alkalic Dry



### Preparation instructions

1. Weigh the required quantity of Liquifine® Alkalic dry (6.3 g per 100 ml suspension) and transfer it to the final recipient<sup>(\*)</sup>.
  2. Weigh the required quantity of omeprazole and transfer it also to the final recipient.
  3. Shake firmly.
  4. Add purified water to the desired volume and shake firmly until a homogeneous dispersion is obtained.
- (\*) You can also use the pre-filled 100 mL or 200 mL bottles with mark line, which contain already the required amount of Liquifine® Alkalic dry.

### Production report

| Preparation date       | / /                      | Galenic type      | Suspension       |
|------------------------|--------------------------|-------------------|------------------|
| Batch number           |                          | Volume            |                  |
| Raw material           | Number analysis register | Required Quantity | Weighed quantity |
| Omeprazole             |                          | g                 | g                |
| Liquifine® Alkalic dry |                          | g                 | g                |
| Purified water         |                          | Ad mL             | Ad mL            |

### Tarra

• \_\_\_\_\_

### Quality control

- Homogeneity: \_\_\_\_\_

### Stability

- 60 days

### Storage and labelling

- Protected from light – Fridge
- Shake before use

| Name and Signature Compounder | Name and Signature Pharmacist |
|-------------------------------|-------------------------------|
|                               |                               |

### MAGIS PHARMA

Keizershoek 336, 2550 Kontich, T: +32 3 457 11 76, F: +32 3 457 43 40  
info@magis-pharma.be, www.magis-pharma.be